

CAUTION

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

Country
(logo)

2,4-D AMINE 500

SELECTIVE HERBICIDE

Active Constituent: 500g/L 2,4-D present as the Dimethylamine Salt

GROUP	I	HERBICIDE
-------	---	-----------

For the control of broadleaved weeds in fallow before direct drilling or sowing of cereals and pastures; and in cereal crops, pastures, sugar cane peanuts and non-agricultural areas as per the Directions for Use table.

READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT.

Manufactured by:
A & C Rural Pty Ltd
A.C.N. 010 865 317
Lot 96 Potassium Street
NARANGBA QLD 4504
Phone: (07) 3203 3944

D.O.M:

Contents: * Litres

Batch No:

NRA Approval No: **50637/0302**

* FOR PACK SIZES 5 L, 20L, 110L, 200L and 1000L

GENERAL INSTRUCTIONS:

Before opening, carefully read Directions for Use, Precautionary Statement, Safety Directions and First Aid Instructions.

APPLICATION INFORMATION:

BOOM SPRAYING - Use 30-120 litres water/ha.

AERIAL SPRAYING - Use 10-90 litres water/ha.

EQUIPMENT MAINTENANCE AND USAGE: Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants, unless it has been well washed out with hot soapy or 1% solution of ammonia, followed by several clear water rinses.

COMPATIBILITY: This product can be tank mixed with Banvel®, Glean®, Siege®, Gramoxone®, Propon™, Nu-trazine®, Spray.seed® Flowable Diuron, and Tribunil®. Always refer to the label of any product being mixed with Country 2,4-D Amine 500 Selective Herbicide, and follow all instructions and restrictions relating to its use.

GROUP	I	HERBICIDE
-------	---	-----------

RESISTANT WEEDS WARNING: Country 2,4-D Amine 500 Selective Herbicide is a member of the Phenoxy group of herbicides and has the Disruptors of plant cell growth mode of action. For weed resistance management, Country 2,4-D Amine 500 is a Group I herbicide.

Some naturally occurring weed biotypes resistance to Country 2,4-D Amine 500 and other disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Country 2,4-D Amine 500 or other disruptors of plant cell growth herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, A & C RURAL PTY LTD Limited accepts no liability for any losses that may result from the failure of Country 2,4-D Amine 500 to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS: DO NOT spray in high winds. DO NOT spray crops or weeds outside the stages indicated in the Critical Comments as damage, loss of yield or inadequate weed control may result. DO NOT spray cereals if lucerne is present.

Drift Warning: DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift onto susceptible crops such as cotton, tobacco, tomatoes, vines, fruit trees, vegetables, legume crops and pastures, oilseed crops and susceptible trees (e.g. Kurrajongs, Belahs and Eucalypts).

PROTECTION OF LIVESTOCK: Low hazard to bees. May be applied on any plants at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT: DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL: Store in the closed, original container in a cool, dry, well-ventilated area away from children, animals and food, feedstuffs, seed and fertilisers. Do not store for prolonged periods in direct sunlight. Triple (or preferably) pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS: Poisonous if swallowed. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray wear PVC or rubber apron, elbow-length PVC gloves and face shield. When using the prepared spray wear face shield. If product on skin immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor, or Poisons Information Centre (Ph: 13 11 26). For further information, refer to the Material Safety Data Sheet (MSDS).

CONDITIONS OF SALE: The use of COUNTRY 2,4-D AMINE 500 being beyond the control of the manufacturer no warranty expressed or implied is given by A & C RURAL PTY LTD regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and A & C RURAL PTY LTD accepts no responsibility for any consequence whatsoever resulting from the use of this product.

COUNTRY**2,4-D AMINE 500
Selective Herbicide**

Active Constituent: 500g/L 2,4-D PRESENT AS THE
DIMETHYLAMINE SALT

GROUP	I	HERBICIDE
--------------	----------	------------------

For the control of broadleaved weeds in fallow before direct drilling or sowing of cereals and pastures; and in cereal crops, pastures, sugar cane, peanuts and non-agricultural areas as per the Directions for Use table.

Manufactured by:
A & C Rural Pty Ltd
A.C.N. 010 865 317
Lot 96 Potassium Street
NARANGBA QLD 4504
Ph: (07) 3203 3944

READ THOROUGHLY BEFORE USE

GENERAL INSTRUCTIONS:

Before opening, carefully read Directions for Use, Precautionary Statement, Safety Directions and First Aid Instructions.

APPLICATION INFORMATION:

BOOM SPRAYING - Use 30-120 litres water/ha.
AERIAL SPRAYING - Use 10-90 litres water/ha.

EQUIPMENT MAINTENANCE AND USAGE:

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants, unless it has been well washed out with hot soapy or 1% solution of ammonia, followed by several clear water rinses.

COMPATIBILITY: This product can be tank mixed with Banvel®, Glean®, Siege®, Gramoxone®, Propon™, Nu-trazine®, Spray.seed® Flowable Diuron, and Tribunil®. Always refer to the label of any product being mixed with Country 2,4-D Amine 500 Selective Herbicide, and follow all instructions and restrictions relating to its use.

GROUP	I	HERBICIDE
--------------	----------	------------------

RESISTANT WEEDS WARNING: Country 2,4-D Amine 500 Selective Herbicide is a member of the Phenoxys group of herbicides and has the Disruptors of plant cell growth mode of action. For weed resistance management, Country 2,4-D Amine 500 is a Group I herbicide.

Some naturally occurring weed biotypes resistance to Country 2,4-D Amine 500 and other disruptors of plant cell growth herbicides may exist through normal genetic

variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Country 2,4-D Amine 500 or other disruptors of plant cell growth herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, A & C RURAL PTY LTD Limited accepts no liability for any losses that may result from the failure of Country 2,4-D Amine 500 to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS:

DO NOT spray in high winds. DO NOT spray crops or weeds outside the stages indicated in the Critical Comments as damage, loss of yield or inadequate weed control may result. DO NOT spray cereals if lucerne is present.

Drift Warning: DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift onto susceptible crops such as cotton, tobacco, tomatoes, vines, fruit trees, vegetables, legume crops and pastures, oilseed crops and susceptible trees (e.g. Kurrajongs, Belahs and Eucalypts).

PROTECTION OF LIVESTOCK: Low hazard to bees. May be applied on any plants at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT: DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL: Store in the closed, original container in a cool, dry, well-ventilated area away from children, animals and food, feedstuffs, seed and fertilisers. Do not store for prolonged periods in direct sunlight. Triple (or preferably) pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS: Poisonous if swallowed. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray wear PVC or rubber apron, elbow-length PVC gloves and face shield. When using the prepared spray wear face shield. If product on skin immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor, or Poisons Information Centre (Ph: 13 11 26)
For further information, refer to the Material Safety Data Sheet (MSDS).

CONDITIONS OF SALE: The use of COUNTRY 2,4-D AMINE 500 being beyond the control of the manufacturer no warranty expressed or implied is given by A & C RURAL PTY LTD regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and A & C RURAL PTY LTD accepts no responsibility for any consequence whatsoever resulting from the use of this product.

N.R.A. Approval No: 50637/0302

IMPORTANT:

**WHEN APPLIED TO DRY SOILS,
AT LEAST 15mm ($\frac{1}{2}$ inch) OF RAIN MUST FALL
PRIOR TO THE COMMENCEMENT OF THE PLANT BACK PERIOD.**

NOTES:

- % In Queensland, no rainfall is required to fall prior to commencement of Plant Back Period for Wheat, Barley and Triticale
- # In Queensland, planting of Canola/Rapeseed, Chickpeas and Safflower must be delayed for at least 14 days following rainfall of at least 15mm.
- @ In Central Queensland, when using 1L/Ha or less of Country 2,4-D Amine 500, the plant back period for Sorghum and Sunflower is 1 day irrespective of rainfall.
- + In W.A. the plant back period for Lupins at all rates is 28 days.

DIRECTIONS FOR USE

RESTRAINTS: DO NOT spray if rain seems likely within 6 hours or if strong winds prevail.
DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

1. FIELD CROPS

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments			
Wheat	Refer Weed Table	N.S.W., S.A. only	400mL - 2.1L	Apply after the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller (N.S.W., S.A. only) Apply from tillering to boot stage (Vic. only) Apply from mid tillering to before boot stage. (Qld. only) Apply at 5 leaf to fully tillered (Tas. only)			
		Vic. only	280mL - 1.7L				
		Qld. only	700mL - 1.7L				
		Tas. only	1.7L				
		Barley	N.S.W., S.A. only		N.S.W., S.A. only	400mL - 1.7L	
					Vic. only	280mL - 1.7L	
					Qld. only	700mL - 1.7L	
					Tas. only	1.7L	
					Cereal Rye, Triticale	N.S.W., S.A. only	N.S.W., S.A. only
		Vic. only	280mL - 1.7L				
Oats	N.S.W., S.A. only	400mL - 1L					
	Vic. only	280mL - 1L					
Cereals: Wheat, Oats, Barley	Cape Tulip	W.A. only	800mL - 1.6L	Apply from the 5 leaf stage up to jointing stage (Zadoks 15-33). Apply after the 6 leaf stage (Z. 16) for Cranbrook, Jacup, Aroona and Spear Wheat and Mortlock Oats to avoid possible damage. DO NOT spray if lucerne is present. WEED STAGE: 10-15cm. Docks should be sprayed before 5 leaf stage. Cape Tulip - low rate for corns only.			
			Dock, Saffron Thistle		1.4L		
			Indian Hedge Mustard, London Rocket, Lupin, Rapistrum, Wild Radish		1L		
			Wild Turnip		800mL		
			Capeweed, Doublegee, Erodium, London Rocket, Lupin, Mustard, Rapistrum, Wild Radish, Wild Turnip		250mL/ha plus 500mL/ha Flowable Diuron 500g/L		
Wheat Barley	Wild Radish	Vic., S.A., N.S.W. only	100mL/ha plus 850g/ha Bayer Tribunil Herbicide	Spray 2-6 weeks after sowing and not later. Do not use on crops undersown with lucerne.			
Fallow, Stubble: Spray prior to Direct Drilling or Sowing a) Winter Cereals b) Winter Cereals and Maize, Sweetcorn, Peanuts Maize, Sweetcorn	Refer Weed Table	Vic. only	280mL - 1.7L	Observe plant back periods given in the table on this leaflet. Can be mixed with Glean®, Gramoxone®, or Spray, Seed® where grasses are present. For Skeleton Weed, spraying should only be done 6-8 weeks before anticipated sowing date and subsequent cultivation limited to a minimum.			
					N.S.W. only	700mL - 2.1L	
					Qld. only		
		Maize, Sweetcorn	N.S.W., S.A. only		700mL - 1.4L	Apply when crop is 10-20cm high and secondary roots are developing for an over the top spray. When crop is between 20cm high and just before tasselling, spray with dropped nozzles to avoid chemical being sprayed into the whorl and on upper leaves.	
					Qld. only	700mL - 1.1L	Apply when crop is 10-30cm high and secondary roots have developed. Use drop nozzles and direct the spray when crop exceeds 30cm in height and before tasselling.
					Tas. only	1.7L	Apply when the crop is 15-30cm high. do not spray if the crop is showing signs of stress. Some leaf twisting may occur following application - crop recovers quickly.
		Sorghum	N.S.W., S.A., Vic. only		700mL - 1.4L	Apply preferably when crop is at 3-6 fully expanded leaf stage but can be sprayed from 2-8 leaf stage. From 6 leaf stage onwards to within 2 weeks of flowering, crop can be sprayed with dropped nozzles to avoid chemical being sprayed into the whorl and on upper leaves.	
					Qld. only (except central Qld)	700mL - 1.1L	Apply when crop has 4 to 8 fully expanded leaves and secondary roots have developed.
		Millet	N.S.W., S.A., Vic. only		700mL - 1.4L	Spray when secondary roots have developed, when fully tillered and before heads start to form at the base of the tillers. Do not use on Panorama Millet or Panicum.	
					Qld. only	700mL - 1.1L	
Maize, Sweetcorn, Sacaline, Broom Millet, Millet	Cape Tulip, Dock, Saffron Thistle, Indian Hedge Mustard, London Rocket, Lupin, Rapistrum, Radish, Wild Turnip	W.A. only	1.4L	Spray when crop is 10-30cm high and secondary roots have developed and before tasselling. Apply as direct spray to weeds.			
			Grain Sorghum	Apply when crop is 12cm high. DO NOT apply between tassel and dough stage. Avoid spraying when in flower.			
Sugar Cane	Bindy Eye (Star Burr), Blue Top, Cobblers Pegs, Fleabanes, Jute, Leucas, Needle Burr, Spear Thistle, Water Primrose, Ipomea Vines, Convolvulus Vines	Qld. only	2.2 - 4.4L	Add 500mL of 60% Non-ionic Surfactant to 100L of spray mixture. Agitate well. DO NOT USE ON Q63, Q67, Q80, OR Q96 VARIETIES.			
			Chinese Mint, Blue Snakeweed		4.4L		
Peanuts	Broadleaf Weeds, except Noogoora Burr, Grasses, except Mossman Burr	Qld. only	2.2L OR 4.5L	LOWER RATE: Apply as BAND SPRAY as soon as possible after planting in a 55cm band. HIGHER RATE: Apply as OVERALL SPRAY after planting and before crop emergence. Some crop damage may occur if heavy rain falls between application and crop emergence.			
Harvest Aid or Salvage Spray - Winter Cereals - Maize & Sorghum	Desiccate Broadleaf Weeds	Qld. only	1.5 - 2.1L	Apply after dough stage.			
	Refer Weed Table	Qld. & N.S.W. only					

2. PASTURES, NON-AGRICULTURAL, RIGHTS OF WAY, INDUSTRIAL LAWNS

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments	
Pastures and Non-agricultural	Refer Weed Table	N.S.W, Qld., S.A., Tas. only	700mL - 2.1L	Pasture legumes including lucerne, clovers and medics may be damaged unless well protected by grasses. Spot spraying is preferred	
			W.A. only	1.4L	For pastures not containing legumes. Only seeding Docks, Spear Thistle and Saffron Thistle will be controlled.
				2L/ha plus 1% crop oil	Spray when plants are actively young preferably before flowering or vining.
				1 - 1.4L	
				2.0L	Spray when plants are small.
				2.0 - 4.0L plus 500ml of 60% Non-ionic surfactant/100Lts of spray mix	Best results are obtained when plants are small. Use high rate on larger plants.
				4.0L	Spray after good emergence of seedlings.

2. PASTURES, NON-AGRICULTURAL, RIGHTS OF WAY, INDUSTRIAL LAWNS (cont)

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Pastures, Rights of Way & Industrial	Boxthorn, Boneseed, Hawthorn	Vic., S.A. only	100mL/10L water	Spot Spraying. For Boneseed only, thoroughly wet plants or seedlings.
	Groundsel	N.S.W., Qld., S.A. only	Undiluted	Apply as an undiluted spray to freshly cut stumps.
			1.2L/15L water	MISTING: Lightly wet plants
			400mL/100L water 300mL/15L water	HIGH VOLUME: Thoroughly wet plants CUT STUMP: Swab the cut stump immediately. Apply by a pouring can or knapsack spray.
	Lantana	N.S.W., Qld., S.A. only	3.6 - 5.5L	AERIAL APPLICATION: Spray when Groundsel is actively growing
			400mL/100L water	Use a coarse spray with sufficient pressure to penetrate canopy and wet stems as well as foliage. Spray at the end of a wet summer (March to May). Defoliation should occur but respraying of new growth will be necessary in following Autumn. Broadcast grass seed and keep stock off following Summer to allow the pasture to establish. Damage may result to pasture legumes.
	Mother of Millions	N.S.W. only	500mL/100L water	Hand gun and knapsack only. A thorough coverage of leaves and plantlets is necessary. Use Spraymate Chemwet 1000 at the rate of 1.0mL of surfactant per 1L of mixture.
Noogoora Burr, Weir Vine (Ipomea)	Qld. only	200mL/100L water	In all cases apply to young, actively growing weeds, ensuring thorough coverage.	
Annual & Perennial Pigweed, Artichoke thistle, Bathurst Burr, Billygoat Weed, Blue snake Weed, Burr Medic, Clockweed*, Fleabanes, Galvanised Burr, Hemlock, Hoary Cress*, Kyalinga Weed (Whisker Grass), Knobweed, Milky Cotton Bushes, Parthenium Weed, Paterson's Curse, Saffron Thistle, Star Burr, Thornapple, Variegated Thistle*		400mL/100L water	* Spray rosette stage - Clockweed, Variegated Thistle ° Repeat spraying necessary - Hoary Cress	
Rubber Vine		200mL/10L water	Apply to freshly cut stump.	
Conservation Tillage - Direct Drilling, Surface Sowing or Fallow Maintenance of Pastures	Charlock, Mustards, Shepherd's Purse, Saffron, Slender, Spear and Variegated Thistles, Turnip Weed, Wild Radish, Wild Turnip	All States	700mL - 2.0L	Apply to actively growing young plants. Before sowing: Observe plant back periods given in the table on this leaflet.
	Clover, Sorrel		1.4L/ha plus 700mL - 1L/ha of Banvel®	Apply to actively growing plants in autumn. Do not sow pasture seed for at least 30 days after application.
Pastures - Spray/Graze Techniques				*PRECAUTION: An increased quantity of poisonous plants may be eaten by stock using Spray-Graze e.g. Catnip, Capeweed, Paterson's Curse and Variegated Thistle and deaths could result from causes such as nitrate poisoning. With Paterson's Curse, preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. Do not graze horses or pigs on Paterson's Curse.
	Amsinckia, Thistles, Capeweed, Doublegee, Mustard, Paterson's Curse, Wild Turnip, Wild Radish, Docks, Geranium, Erodium	S.A. only	700mL	Apply from 6 weeks after opening rains in autumn until the end of August. Seven days after spraying, stock paddock at 4-5 times normal rate, preferably with sheep. Maintain this level of grazing for 6 weeks or until pasture shows signs of over grazing. Then return to normal stocking levels. Use high stocking rates in following spring to prevent weeds from flowering. Repeat treatments may be required for 2-3 years for complete control.
	Annual Thistles, Capeweed, Doublegee, Mustards, Paterson's Curse, Turnip, Saffron Thistle, Spear Thistle	Tas, Vic. only		
	Amsinckia, Docks (seedling only), Capeweed, Doublegee, Mustard, Wild Radish, Wild Turnip, Paterson's Curse, Annual Thistles	W.A. only	750mL	
	Spear Thistle, Saffron Thistle		1.5L	Apply to Saffron Thistle at the end of September when plants are running up to flower. Sub-clovers may be damaged at this rate and use is not recommended for all medic pastures.
	Melons		2.0L + 1% oil	Heavy stocking on young plants sprayed with 1.0L/ha provides effective control.
	Docks	Vic. only	1.4L	Apply in September only. Sub-clovers may be damaged at this rate and use is not recommended for all medic pastures.
Callitriche, Capeweed, Charlock, Mustards, Paterson's Curse, Shepherd's Purse, Saffron, Slender, Spear or Variegated Thistle, Turnip Weed, Wild Radish, Wild Turnip	N.S.W. only	350mL - 1.4L	Spray actively growing 6-8 week old weeds. Introduce stock 7-10 days after spraying, preferably sheep (cattle are less effective). Stocking rate should be at least 5 times heavier than normal until weeds have been reduced, but before survival of desirable pasture species is threatened. Lucerne and Medics may be damaged and should be grazed short before spraying. Other legumes may be affected.	
Lawns	Refer Weed Table	Qld. only	2-4mL/1L water	Wet foliage thoroughly.
		W.A. only	50mL/100mL	

3. SPOT SPRAYING

Situation	Weeds Controlled	State	Mixing Rates/Comments
High Volume Spraying	Refer to Weed Table	All States	Add 1/10th of rate on weed table to 150 litres of water. Each 150 litres of mix will cover 1000m ² (1/10th Ha). e.g. if rate in weed table is 1.4L, use 140mL/150L water.
Knapsack Application	Refer to Weed Table	All States	Add 1/100th of rate on weed table to 10 litres of water. Each 10 litres of mix will cover 100m ² (1/100th Ha). e.g. if rate in weed table is 1.4L use 14mL/10L water.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: PASTURE, CEREAL CROPS - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

IN TASMANIA, THIS PRODUCT MAY ONLY BE USED FROM 15 APRIL TO 15 SEPTEMBER UNLESS OTHERWISE PERMITTED BY THE REGISTRAR OF PESTICIDES.

WEED TABLE:

NOTE: Where weeds are to be sprayed in a CROP or PASTURE, use only the rates given for the particular crop or situation indicated under the Directions for Use Table. In most cases, this will give control, however, some hard to kill weeds or those in advanced stages of growth may only be suppressed.

The rates listed in the Weed Table below are for use where weeds only are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage.

This product should only be used in those stages where a rate is indicated for the particular weeds listed.

WEED TABLE:

Weeds Controlled	Vic.	APPLICATION RATE PER HECTARE						CRITICAL COMMENTS
		CROP WEEDS			PASTURE WEEDS			
		N.S.W.	S.A.	Qld.	Tas	W.A.	N.S.W., S.A., Qld., Tas only	
Amaranthus spp.	-	700mL-1.4L	-	1.1L	-	-	-	Spray young plants
Apple of Peru	-	700mL-1.4L	-	1.1L	-	-	-	Spray young plants. Susceptible when young.
Bathurst Burr	-	1.4L	-	1.1L	-	-	1-1.4L NOT S.A.	Spray seedlings only
Blackberry Nightshade	-	700mL-1.4L	-	1.1L	-	-	-	-
California Burr	-	1-1.4L	-	1.1L	-	-	1-1.4L NOT S.A.	Spray seedlings only
Cape Tulip	-	-	-	-	-	800mL-1.6L	-	Low rate for cormils only.
Capeweed	1.7L	-	2.1L	-	1.7L	-	2.1-3.5L	Spray seedlings to rosette stage
Caltrop	-	1-2.1L	-	1.1L	-	-	-	Moderately susceptible
Charlock	700mL-1L	700mL-1.4L	700mL	-	1.7L	-	1-1.4L	Spray at rosette stage.
Clover	-	1.5L	-	-	-	-	-	-
Common ice Plant	-	-	1.4L	-	-	-	-	-
Docks	1.7L	-	1.7L	1.7L	1.7L	1.4L	4L S.A. ONLY	Spray at multiple leaf stage -effective only on seedlings.
Fat Hen	-	700mL-2.1L	-	1.1L	1.7L	-	-	Spray pre-flowering
Fumitory - red	-	-	2.1L	-	-	-	-	-
Fumitory - white	1L	-	700mL	-	-	-	-	Spray at multiple leaf stage.
Hexham Scent/Mellilotus	1.7L	-	1.4L	1.7L	-	-	1.4-2.1L	Spray multiple leaf stage before seeding.
Hoary Cress	1.1-1.7L	1.4-2.1L	1.7L	1.7L	-	-	1.8-2.1L	Spray rosettes and pre-flowering.
Hogweed/Wireweed	1.7L	-	-	1.7L	-	-	-	Spray at multiple leaf stage (Vic.) Spray at seedling. spray at seedling and young plant stage (Qld).
Horehound	-	-	1.7L	-	-	-	2.8-4L S.A. only	Spray seedlings.
Khaki Weed	-	-	-	-	-	-	1.4-2.8L Not S.A	Spray seedlings only
Lincoln Weed	-	-	2.1L	-	-	-	-	Spray early rosettes.
London Rocket	-	-	-	-	-	1.0L	-	-
Lupins	-	1-2.1L	-	-	-	-	-	-
Mexican Poppy	-	-	-	1.7L	-	-	-	Spray seedlings - plants become more resistant with age.
Mintweed	-	1.4L	-	1.1L	-	-	-	Spray seedlings - resistance in later stages
Mustards	280-700mL	700mL-1.4L	700mL-1.7L	1.1L	-	1.0L	700mL-1.4L	Spray at 2-4 leaf up to rosette stage.
New Zealand Spinach	-	1.4L-2.1L	-	-	-	-	-	-
Noogoora Burr	-	1-1.4L	-	1.1L	-	-	1-1.4L NOT S.A.	Spray seedlings only.
Paterson's Curse	-	1.4-2.1L	-	1.7L	-	1.6L	2.1-2.8L	Spray rosettes or before plants have 10 leaves. Later stages harder to kill.
Potato Weed	-	700mL-1.4L	-	1.1L	-	-	-	-
Rapesed	-	1-2.1L	-	-	-	-	-	-
Rough Poppy	-	1.4L	-	-	-	-	-	-
Safflower	-	700mL-1.4L	-	-	-	-	-	-
Shepherd's Purse	-	1.4-2.1L	-	-	1.7L	-	1-1.4L	Spray young rosettes.
Skeleton Weed	1.7L	1.4-2.1L	1.7L	-	-	-	1.8-2.8L	Spray rosettes before aerial growth commences.
Sorrel	1.7L	2.1L	1.7L	-	-	-	-	Only moderately susceptible.
Speedwell-ivy Leaf	-	-	1.4L	-	-	-	-	-
Spiny Emex	-	-	-	1.7L	-	-	-	Only young plants are susceptible
Stinkwort	-	1-1.7L	-	-	-	-	-	-
Storkbill/Erodium	-	-	-	-	1.7L	-	2-4L	Spray seedlings to young rosettes.
Sunflower(seedlings)	1.7L	700mL-1.7L	-	1.1L	-	-	-	-
Thistle	-	-	-	-	-	-	-	-
- Californian	-	-	-	-	3.4L	-	4-4.7L	Repeated applications may be necessary. (N.S.W., Tas only).
- Saffron	1.4L	700mL-2.1L	1.7L	1.7L	1.2L	1.4L	1.4-2.1L	Low rate only sufficient to control weeds in crops at rosette stage when spot-sprayed early.
- Slender/Shore	-	1-2.1L	-	-	1.7L	-	1.4L	Suppression only
- Soldier	1.7	-	-	-	-	-	1.4-2L	Spray young rosette
- Spear	700mL	-	-	-	1.7L	-	1.4-2.1L	Spray young rosettes
- Star	-	-	-	-	-	-	2-4L S.A. ONLY	Use higher rate as flower stalk appears.
- Variegated	-	700mL-2.1L	-	1.1L	1.7L	-	1.4-2.1L	Spray at rosette stage
Thornapple	-	1-1.4L	-	-	-	-	2-3L NOT S.A.	Spray seedlings only.
Turnip Weed/Rapistrum	-	700mL-1.4L	-	700mL	-	1.0L	700mL-1.4L	-
Wards Weed	-	-	1.4L	-	-	-	-	-
Wild Cabbage	1.7L	-	-	-	-	-	-	Spray multiple leaves
Wild Poppy	700mL	-	-	-	-	-	1.4-2.1L	Spray rosettes
Wild Radish	1.7L	1.7-2.1L	1.7L	1.1L	1.7L	1.0L	1-1.4L	Spray up to young rosette stage
Wild Turnip	280-700mL	700mL-1.4L	400mL	-	1.7L	800mL	700mL-1.4L	Spray 2-4 leaf up to rosette stage
Vetches/Tares	1.7L	-	1.4L	-	-	-	-	Spray at multiple leaf stage.

PLANT BACK FOR COUNTRY 2,4-D AMINE 500

CROP	RATES:		
	Up to 700mL/Ha	700mL-1.4L/Ha	1.4L - 2.1L/Ha
Balansa Clover	7	7	10
Barley %	1	1	3
Chickpeas#	7	14	21
Cotton	10	14	21
Faba Beans	7	7	10
Field Peas	7	14	14
Lentils	7	7	10
Linseed	7	7	14
Lucerne	7	7	10
Lupins+	7	14	21
Medic	7	7	10
Narbon Beans	7	7	10
Navybean	10	10	14
Oats	3	3	7
Perennial Ryegrass	7	7	10
Persian Clover	7	7	10
Phalaris	7	7	10
Canola/Rapeseed #	14	21	28
Rice	7	7	14
Safflower #	7	14	21
Sorghum @	3	7	10
Soybean	14	14	21
Sub-Clover	7	7	10
Sunflower @	7	10	14
Triticale %	1	3	7
Vetch	7	7	10
Wheat %	1	3	7
White Clover	7	7	10