## **POISON** KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



## MICRO-LO PRE-CONSTRUCTION/POST-CONSTRUCTION TERMITICIDE AND INSECTICIDE

ACTIVE CONSTITUENT:450 g/L CHLORPYRIFOS (an anticholinesterase compound)SOLVENT:318.5g/L LIQUID HYDROCARBON



For installing chemical soil barriers for new building work in accord with AS 3660.1 and for the post-construction management of subterranean termites in accord with AS 3660.2 and other domestic insect pests as specified in the Directions for Use table.

> IMPORTANT: RESTRICTED CHEMICAL PRODUCT ONLY TO BE SUPPLIED TO, OR USED BY, AN AUTHORISED PERSON

THIS PRODUCT IS TOO HAZARDOUS FOR USE BY HOUSEHOLDERS. HOUSEHOLDERS MUST NOT USE THIS PRODUCT IN OR AROUND THE HOME.

#### IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE

A & C RURAL PTY LTD ACN 52 010 865 317 Lot 96 Potassium Street NARANGBA QLD 4503 Ph: (07) 3897 2000

D.O.M.: Batch no: NRA Approval Nº: 53192/0103

Contents: FOR PACK SIZES 5LT, 20LT, 110LT, 200LT & 1000LT

#### STORAGE AND DISPOSAL

Store in closed, original container in a cool well-ventilated area. Do not store for long periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling break, crush or puncture and bury empty containers in a local authority landfill If no local landfill is available, bury the containers at a depth of 500mm or deeper in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. Rinsate/rinse water should be disposed of in accordance with appropriate State legislation and should preferably be sprayed on to an application site or added as up to 10% of the diluent the next time this product or another similar termiticide is used. Do not put down sewers, gutters or storm water drains. In some States wastes can only be buried at a licensed landfill.

#### SAFETY DIRECTIONS

#### <u>For General Use</u>

Product is poisonous if absorbed by skin contact, inhaled or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Obtain an emergency supply of atropine tablets 0.6 mg. Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale vapour or spray mist.

When opening the container, preparing the spray or using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield or goggles.

If product is spilt on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

#### For Use As A Termiticide

Product is poisonous if absorbed by skin contact, inhaled or swallowed. Repeated exposure may cause allergic disorders. Repeated minor exposure may have a cumulative poisoning effect. Obtain an emergency supply of atropine tablets 0.6 mg. Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale vapour or spray mist.

When opening the container, preparing the spray and using the prepared spray, wear chemical resistant clothing buttoned to the neck and wrist and washable hat, elbow-length PVC gloves, goggles, chemical resistant footwear and half facepiece respirator with combined dust and gas cartridge.

If clothing becomes contaminated with product or wet with spray, remove clothing immediately. If product on the skin, immediately wash area with soap and water. If product is in the eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, goggles, respirator (and if rubber wash with detergent and warm water), and contaminated clothing.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. (Ph:. 13 11 26). If swallowed, give one atropine tablet every 5 minutes until dryness of the mouth occurs - if poisoned by skin absorption or through lungs, remove any contaminated clothing, wash skin thoroughly and give atropine tablets as above. Get to a doctor or hospital quickly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

#### MATERIAL SAFETY DATA SHEET

Additional information is listed on the Material Safety Data Sheet for Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide which is available from A & C Rural Pty Ltd on request.

**CONDITIONS OF SALE:** The use of Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide being beyond the control of the manufacturer no warranty expressed or implied is given by A & C Rural Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and A & C Rural Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.

Page 2 of 16	27/04/05
Country Micro-Lo Pre-Construction/Post C	onstruction Termiticide & Insecticide

UN NO. 3018 ORGANOPHOSPHORUS, PESTICIDE, LIQUID, TOXIC. PKG III HAZCHEM 2X



### **POISON** KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# MICRO-LO PRE-CONSTRUCTION/POST-CONSTRUCTION TERMITICIDE AND INSECTICIDE

ACTIVE CONSTITUENT:450 g/L CHLORPYRIFOS (an anticholinesterase compound)SOLVENT:318.5g/L LIQUID HYDROCARBON

# GROUP 1B INSECTICIDE

For installing chemical soil barriers for new building work in accord with AS 3660.1 and for the post-construction management of subterranean termites in accord with AS 3660.2 and other domestic insect pests as specified in the Directions for Use table.

> IMPORTANT: RESTRICTED CHEMICAL PRODUCT ONLY TO BE SUPPLIED TO, OR USED BY, AN AUTHORISED PERSON

THIS PRODUCT IS TOO HAZARDOUS FOR USE BY HOUSEHOLDERS. HOUSEHOLDERS MUST NOT USE THIS PRODUCT IN OR AROUND THE HOME.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE

A & C RURAL PTY LTD ACN 52 010 865 317 Lot 96 Potassium Street NARANGBA QLD 4503 Ph: (07) 3897 2000

NRA Approval no: 53192/0103

#### **DIRECTIONS FOR USE:**

#### MANAGEMENT OF SUBTERRANEAN TERMITES (All States, except Tasmania) RESTRAINTS

DO NOT apply to soils if excessively wet, immediately after heavy rain or if heavy rains are expected within 48 hours to avoid run-off of chemical.

**DO NOT** use at less than indicated label rates.

DO NOT use in cavity walls, except for direct treatment of nest.

DO NOT use on alkaline soils in SA (use on neutral or acid soils only) or on dolomite based sub-slab bedding material.

#### **RATES OF APPLICATION**

**IMPORTANT:** Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide should be used as part of an overall termite management program as detailed in Australian Standard Series AS 3660. A great deal of care is required to understand construction details of the building and to apply the product in a manner which ensures a complete chemical soil barrier. Where necessary, the barrier may need to be re-applied under the building. Application equipment must be fitted with a flow meter and pressure regulator on the application device. The purpose of a chemical soil barrier is to impede and discourage concealed termite entry into a structure. Barriers may still be bridged by termites, but their entry can then be more easily detected during routine inspections. If a barrier is not complete or breached, then concealed termite entry may occur. It is often not possible to form a complete barrier around existing structures in which case other termite management options and/or more frequent inspections will also need to be considered.

#### **PRE-CONSTRUCTION USAGE**

Page 5 of 16

Installation of chemical soil barriers prior to of houses, factories, industrial and commercial buildings, public premises, and farm buildings.Subterranean termites, south of the Tropic of CapricornHorizontal Barriers: Use 110 mL/m2Horizontal Barriers: Use 110 mL/m2Use 110 mL of Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide/5L of water and apply mixture (the emulsion) at 5L/m2. See APPLICATION VOLUME in GENERAL INSTRUCTIONS section for other recomme dilution rates.Vertical Barriers: Use of houses, factories, industrial and farm buildings.Vertical Barriers: Use 2.2L/m3Vertical Barriers: Use 2.2 L of Country Micro-Lo Pre-Construction/Post-Construction Post-Construction/Post-Construction Termiticide and Insecticide/100L of water and apply at 100L mixture per m3 soil. Dilution rate: 220 mL/10L. 220 mL/10L is equivalent to a 1% active ingredient emulsion. See Service requirement in GENERAL INSTRUCTIONS for expected barrier life.SubterraneanHorizontal Barriers:Horizontal Barriers:Horizontal Barriers:	
completion construction of houses, factories, industrial and commercial buildings, public premises, and farm buildings.the Tropic of CapricornAPPLICATION VOLUME in GENERAL INSTRUCTIONS section for other recomme dilution rates.Vertical Barriers: Use 2.2L/m³2.2L/m³Vertical Barriers: Use 2.2 L of Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide/100L of water and apply at 100L mixture per m³ soil. Dilution rate: 220 mL/10L is equivalent to a 1% active ingredient emulsion. See Service requirement in GENERAL INSTRUCTIONS for expected barrier life.	nended
of houses, factories, industrial and commercial buildings, public premises, and farm buildings. Capricorn Vertical Barriers: Use 2.2 L of Country Micro-Lo Pre-Construction/Post-Construction   Public premises, and farm buildings. 2.2L/m <sup>3</sup> Vertical Barriers: Use 2.2 L of Country Micro-Lo Pre-Construction/Post-Construction   Public premises, and farm buildings. 2.2L/m <sup>3</sup> Vertical Barriers: Use 2.2 L of Country Micro-Lo Pre-Construction/Post-Construction   Public premises, and farm buildings. 2.2L/m <sup>3</sup> Vertical Barriers: Use 2.2 L of Country Micro-Lo Pre-Construction/Post-Construction   Public premises, and farm buildings. 2.2L/m <sup>3</sup> Vertical Barriers: Use 2.2 L of Country Micro-Lo Pre-Construction/Post-Construction   Public premises, and farm buildings. 2.2L/m <sup>3</sup> Vertical Barriers: Use 2.2 L of Country Micro-Lo Pre-Construction/Post-Construction   Public premises, and farm buildings. 2.2L/m <sup>3</sup> Vertical Barriers: Use 2.2 L of Country Micro-Lo Pre-Construction/Post-Construction   Public premises, and farm buildings. See Service requirement to a 1% active ingredient emulsion.   See Service requirement in GENERAL INSTRUCTIONS for expected barrier life.	nended
industrial and commercial buildings, public premises, and farm buildings.	
commercial buildings, 2.2L/m³ Vertical Barriers: Use 2.2 L of Country Micro-Lo Pre-Construction/Post-Construction   public premises, and farm buildings. 2.2L/m³ Vertical Barriers: Use 2.2 L of Country Micro-Lo Pre-Construction/Post-Construction   farm buildings. Dilution rate: 220 mL/10L. Dilution rate: 220 mL/10L.   See Service requirement in GENERAL INSTRUCTIONS for expected barrier life. See Service requirement in GENERAL INSTRUCTIONS for expected barrier life.	
public premises, and farm buildings. Termiticide and Insecticide/100L of water and apply at 100L mixture per m³ soil.   Dilution rate: 220 mL/10L. Dilution rate: 220 mL/10L.   220 mL/10L is equivalent to a 1% active ingredient emulsion. See Service requirement in GENERAL INSTRUCTIONS for expected barrier life.	
farm buildings. Dilution rate: 220 mL/10L.   220 mL/10L is equivalent to a 1% active ingredient emulsion.   See Service requirement in GENERAL INSTRUCTIONS for expected barrier life.	
220 mL/10L is equivalent to a 1% active ingredient emulsion. See Service requirement in GENERAL INSTRUCTIONS for expected barrier life.	
See Service requirement in GENERAL INSTRUCTIONS for expected barrier life.	
Subterranean Harizantal Barriars: Harizantal Barriars: (Soa abaya)	
Subtration Horizontal Darriers. Horizontal Darriers. (See above)	
termites north of Use 110 mL/m <sup>2</sup> Vertical Barriers: (See above)	
the Tropic of At this use rate, the expected life of the under-slab treatment is 4 years or more. The expected	ed life
Capricorn or of the external barrier is 2 years or more. This rate must be used in conjunction with a reticul	ulation
where Vertical Barriers: system which is able to apply Country Micro-Lo Pre-Construction/Post-Construction Termit	iticide
Mastotermes Use $2.2L/m^3$ and Insecticide at the required application rates to the required area (ie as per the label direct	ctions
<i>darwiniensis</i> is of for horizontal and vertical barriers and AS 3660 Series), such as Termguard <sup>®</sup> .	
concern. Horizontal Barriers: Horizontal Barriers: Use 220 mL of Country Micro-Lo Pre-Construction/Post-Construction	on
Use 220 mL/m <sup>2</sup> Termiticide and Insecticide/5L of water and apply mixture (the emulsion) at 5L/m <sup>2</sup> . See	
APPLICATION VOLUME in GENERAL INSTRUCTIONS section for other recomme	lended
Vertical Barriers: Use dilution rates.	
4.4L/m <sup>3</sup> Vertical Barriers: Use 4.4L of Country Micro-Lo Pre-Construction/Post-Construction	
Termiticide and Insecticide/100L of water and apply at 100L mixture per m <sup>3</sup> soil.	
Dilution rate: 440 mL/10L.	
440 mL/10L is equivalent to a 2% active ingredient emulsion.	
See Service requirement in GENERAL INSTRUCTIONS for expected barrier life.	

### <u>NOTE</u>: THE SOIL BARRIER PROVIDED BY THIS PRODUCT HAS A FINITE LIFE. THIS, TOGETHER WITH THE RECOMMENDATION TO UNDERTAKE ANNUAL INSPECTIONS, MUST BE STATED ON THE DURABLE NOTICE REQUIRED BY THE B.C.A. at B.1.3.(j)(ii).

APPROPRIATE LEGISLATION

# **APPLICATION INSTRUCTIONS – PRE-CONSTRUCTION CONCRETE SLAB:**

Page 6 of 16

- a) Hand Spray Treatment Before Laying the Slab: The site should be prepared in accordance with Australian Standard AS 2870, or Australian Standard AS 3660.1, as appropriate. Where loose or porous sands occur the site should be moistened with water immediately prior to treatment with Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide, as this will help prevent loss of emulsion through 'piping' or excessive percolation. It will also assist with lateral dispersion. Use a shrouded rose head shower nozzle operated at 170 kPa to apply the required rate of Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide in a volume of 5L of emulsion/m<sup>2</sup>. To obtain better coverage in difficult soil types, it may be necessary to adjust the volume being applied (see APPLICATION VOLUME\_Section under GENERAL INSTRUCTIONS). Ensure application is made under conditions which prevent off-target drift. Immediately following treatment (i.e. on the same day) the moisture membrane should be placed over the treated soil to prevent disturbance of the termiticidal barrier.
- b) External Barrier: On completion of the building, an external barrier should be installed around the perimeter. The external barrier should circumference all pipes and service facilities. External barriers should be created by using either a vertical or horizontal barrier, as determined by the building construction type and adjoining ground level. An external barrier is an essential part of the treatment when relying on a chemical soil barrier to provide the full termite management system.

An external horizontal barrier only, is required when prevention of concealed vertical access by termites is necessary at the perimeter (eg, when ground level is equal to the top of a slab, where the slab is also a barrier to concealed termite movement into the building). A vertical barrier is required when prevention of concealed horizontal access is necessary (eg, where ground level is higher than building material vulnerable to concealed horizontal entry by termites).

- i) Horizontal Barrier: Use a rose head shower nozzle operated at 170 kPa to apply the required rate of 1.5L of the correctly diluted Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide per lineal metre (150mm wide) to soil loosened to a depth of at least 80mm (see APPLICATION VOLUME Section).
- ii) Vertical Barrier: The vertical barrier should be at least 150mm wide and should reach down to 50mm below the top of the footings. To achieve this trench to the top of the footings, and where this is not possible, a combination of trenching (preferably at least 300mm deep) and rodding into the base of the trench may be necessary.

Apply Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide emulsion at 100L per cubic metre of backfill soil, this equates to 1.5L of emulsion/linear metre of a trench 150mm wide and 100mm deep. Where the required vertical barrier is deeper than 100mm, ensure the same rate of application for the extra volume of soil. Use a rose head shower nozzle operated at 170 kPa to flood the base of the open trench and also to treat the backfill soil as it is replaced into the trench to ensure even distribution. Where rodding is necessary, rod before the trench is treated using the spacings in the following table.

#### **Rod Spacings:**

Heavy Clay	ry Clay Clay Loams Sands	
150 mm	200 mm	300 mm

Use a soil injector with a B&G or similar multi-directional tip operated at 170 kPa and insert the rod to the foundation foot as close as possible to the foundation wall ensuring the chemical is applied during insertion and withdrawal. The 4 way tip needs to be rotated during the application through 90 degrees and the 3 way tip through 120 degrees.

#### SUSPENDED FLOORS

Install horizontal and vertical barriers as specified in AS 3660.1 to abut all substructure walls, stumps, piers, pipes and wastes using the techniques described above for <u>external barriers around concrete slabs</u>. Apply the required rate of Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide in a total volume of 5L of emulsion/m<sup>2</sup> (See APPLICATION VOLUME Section). If the barrier is disturbed by any means, retreat to restore continuity. Where the subfloor crawl space is less than 400mm the soil barrier must cover all areas of the subfloor soil.

Note: It is unnecessary to treat the entire subfloor soil where there is access for maintenance and inspection.

SITUATION	RATE	CRITICAL COMMENTS	
Installing a chemical soil		Horizontal Barriers: Use 110mL of Country Micro-Lo Pre-Construction/Post-Construction Termiticide and	
barrier around and under	Barriers:	Insecticide per 5L of water and apply the mixture (emulsion) at a rate of $5L/m^2$ .	
buildings. 1	$10 \text{mL/m}^2$	Vertical Barriers: Use 2.2L of Country Micro-Lo Pre-Construction/Post-Construction Termiticide and	
	Vertical	Insecticide per 100L of water and apply the mixture at a rate of 100L/m <sup>3</sup> .	
	Barriers:	See APPLICATION VOLUME section in GENERAL INSTRUCTIONS for further information.	
2	$2.2L/m^3$	2.2L/100L is equivalent to a 1% active ingredient emulsion.	
		See Service requirement in GENERAL INSTRUCTIONS for expected barrier life.	
Installing a treated soil <b>Horizontal</b> barrier around buildings <b>Barriers:</b> by use through110mL/m <sup>2</sup> Termguard reticulation systems or systems that meet AS Series 3660.		The reticulation system used must be able to apply Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide at the required application rates to the required area (ie. as per the above label directions and AS Series 3660). <b>Termguard:</b> Use 110mL of Country Micro-Lo Pre-Construction/Post-Construction Termiticide and	
		Insecticide/5L of water and apply the mixture at $5L/m^2$ .	
		See APPLICATION EQUIPMENT section in GENERAL INSTRUCTIONS for further information.	
Treatment of termite nest lor colony.		Once the nest or colony has been located it should be broken open and flooded with emulsion. This includes nests located in trees. When treating trees, the addition of a wetting agent is suggested.	
		Refer to Australian Standard Series AS 3660.	

#### **POST-CONSTRUCTION USAGE RATES**

Page 8 of 16	27/04/05
	Country Micro-Lo Pre-Construction/Post Construction Termiticide & Insecticide

Installing a chemical soil	220mL/10L of Trench (preferred) or rod and puddle-treat backfill, ensuring a complete and continuous treated soil barrier is	
barrier around new and	water or creosote provided around the pole or post, to a minimum depth of 300mm and minimum width of 150mm. Use 100L	
existing poles, eg.	of emulsion per m <sup>3</sup> of soil. In addition, infested poles may be drilled near ground level and the cavity flooded	
transmission and building	with the emulsion. This allows seepage to form a treated soil barrier.	
poles, fence posts and	Note: A 50mm gap between fence palings and soil will reduce termite attack and fungal decay. Only soil in	
palings.	contact with palings should be treated.	
_	• Replenishment is recommended within 2 years north of the Tropic of Capricorn and 5 years in other areas.	
	• If the barrier is disturbed, or rain falls immediately after application, retreat to restore continuity and	
	completeness of the barrier.	
	Refer to Australian Standard Series AS 3660.	

®Trademark of Termguard Pty Limited

#### NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

#### **APPLICATION INSTRUCTIONS – POST-CONSTRUCTION**

#### **APPLICATION EQUIPMENT**

#### Hand Spraying

For hand spraying use a rose head shrouded nozzle, operating at 170 kPa, with a flow meter and pressure regulator fitted to the hand-piece.

#### **Treatment Beneath Concrete Slabs or Sealed Areas**

Where it is not possible or practical to remove the slab to allow direct application to the soil, use a sub-slab injector fitted with multi-directional tip (eg. a B&G or similar system) with a 5 degree upward angle (eg. 3 way or 4 way) operated at 170 kPa. Ensure a strong seal with the top of the drill hole to avoid leakage. For the best distribution, the injector needs to be held vertically, at right angles to the slab, and rotated during the application through 90 degrees (if using a 4 way dispersion tip), or through 120 degrees (for a 3 way dispersion tip).

#### **Injection into Soil**

Where it is not possible or practicable to trench the soil; use a soil rod with a 3 or 4 way multi-directional tip (B&G, or similar) operated at 170 kPa. The 4 way tip needs to be rotated during the application through 90 degrees and the 3 way tip through 120 degrees.

#### **APPLICATION VOLUME**

To compensate for impervious soils such as heavy clay where application of  $5L/m^2$  would cause run-off, it may be necessary to apply a volume of emulsion less than  $5L/m^2$ . When reducing the total volume of emulsion used, increase the concentration accordingly to match the label rate by mixing the

required amount of Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide per  $m^2$  in a lesser volume of water. **DO NOT** use emulsion volumes less than 2L for every square metre to be treated.

Note: Use of emulsion volumes other than the recommended  $5L/m^2$  is only permitted when installing barriers in exposed soil. It is not permitted when injecting through the slab or into sealed areas.

#### **Existing Structures**

#### a) Strategic Drilling Through Slab, or Sealed Areas

For treatment of slabs when termites are entering the building through the slab, where reticulation systems do not exist, slab drilling and injection will be required. In most cases, unless there is a known severe termite hazard, grid drilling of the slab is not required. Any such need is to be determined by a licensed Pest Manager.

Treatment needs to be made around the inside of all exterior walls to complete a termite barrier, along both sides of interior wall partitions, around plumbing/electrical or piping entry points and along major cracks or expansion joints. When treating along major cracks or expansion joints it is recommended that holes are drilled alternately on either side of the crack at the recommended drill hole spacings.

For a sand base or sandy soil, apply through a row of holes drilled no more than 300mm apart and 100-200mm out from the wall, crack or pipe. For a clay base, apply through a row of holes drilled 150mm apart and 100mm from the wall, crack or pipe. Apply 10L of emulsion per linear metre and ensure the holes are securely plugged after treatment.

#### b) External Barriers

An external barrier should be installed around the perimeter of the building and should circumference all pipes and service facilities. External barriers should be created by using either a vertical or horizontal barrier, as determined by the building construction type and adjoining ground level. <u>An</u> external barrier is an essential part of the treatment when relying on a chemical soil barrier to provide the full termite management system as per AS 3660.

An external horizontal barrier is only required when prevention of concealed vertical access by termites is necessary at the perimeter (eg, when ground level is equal to the top of a slab, where the slab is also a barrier to concealed termite movement into the building). A vertical barrier is required when prevention of concealed horizontal access is necessary (eg, where ground level is higher than building material vulnerable to concealed horizontal entry by termites).

- i) Horizontal Barrier: Use a rose head shower nozzle operated at 170 kPa to apply the required rate of 1.5L of the correctly diluted Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide per lineal metre (150mm wide) to soil loosened to a depth of at least 80mm (see APPLICATION VOLUME Section).
- ii) Vertical Barrier: The vertical barrier should be at least 150mm wide and should reach down to 50mm below the top of the footings. To achieve this trench to the top of the footings, and where this is not possible, a combination of trenching (preferably at least 300mm deep) and rodding into the base of the trench may be necessary.

Apply Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide emulsion at 100L per cubic metre of backfill soil, this equates to 1.5L of emulsion/linear metre of a trench 150mm wide and 100mm deep. Where the required vertical barrier is deeper than 100mm,

Page 10 of 16	27/04/05
-	Country Micro-Lo Pre-Construction/Post Construction Termiticide & Insecticide

ensure the same rate of application for the extra volume of soil. Use a rose head shower nozzle operated at 170 kPa to flood the base of the open trench and also to treat the backfill soil as it is replaced into the trench to ensure even distribution. Where rodding is necessary, rod before the trench is treated using the spacings in the following table.

#### **Rod Spacings:**

Heavy Clay	Clay Loams	Sands	
150 mm	200 mm	300 mm	

Insert the rod to the foundation foot as close as possible to the house wall ensuring the chemical is applied during insertion and withdrawal. (See **APPLICATION EQUIPMENT Section, Injection into Soil**)

#### c) <u>Suspended Floors</u>

Install horizontal and vertical barriers as specified in Australian Standard Series AS 3660 to adjoin all substructure walls, stumps, piers, pipes and wastes using the techniques described for <u>external barriers around concrete slabs</u>. (See Existing Structures Section.)

### **GENERAL INSTRUCTIONS – Termite Management**

#### CONDITIONS FOR USE BY AUTHORISED PERSONS

The applicator (Pest Manager) must be licensed under State legislation. The Pest Manager must supervise the installation of the moisture membrane and the building workers must wear appropriate protective equipment while laying the moisture membrane.

#### **Termite Management**

To minimise the risk of termite infestation, the subfloor area of buildings should be kept free of stored or waste timber and all other building materials that attract termites. Appropriate action should also be taken to eliminate any undue dampness caused by leaking water or sewerage pipes, or inadequate drainage. Subterranean termites need a constant source of moisture to survive. Provision of adequate ventilation in the subfloor area also helps eliminate undue dampness. Pest managers using this product for termite management should advise either the builder, contractor or home owner, etc, that disturbing the treated soil barrier with subsequent construction of additions or alterations, paths, steps, landscaping, etc, may render the termite management system in place ineffective unless further management options are considered.

#### Colonies not in contact with the ground

Occasionally subterranean termites establish a colony in a building without having contact with the soil because they have access to a continuous supply of moisture (eg. from a faulty plumbing fixture or leaking roof). Such colonies are not affected by chemical soil barriers and should be treated as recommended for established colonies, as per Australian Standard Series AS 3660. Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide may be applied directly to the termite colony in such situations.

#### Service requirement

Page 12 of 16

Regular, competent inspections by a licensed Pest Manager are recommended as part of an overall termite management program to determine the prevailing termite pressure and environmental conditions and consequent requirement for further termite management options. Inspections should be performed at least on an annual basis, but more frequent inspections are strongly recommended.

At the 1% application rate, Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide can provide an effective chemical soil barrier in <u>subfloor regions</u> for 4 years or more north of the Tropic of Capricorn, and for 10 years or more south of the Tropic of Capricorn. At the 1% application rate, Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide can provide an effective chemical soil barrier in <u>exposed situations</u> for 2 years or more north of the Tropic of Capricorn, and up to 5 years or more south of the Tropic of Capricorn.

At the 2% application rate north of the Tropic of Capricorn, Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide can provide an effective chemical soil barrier in <u>subfloor regions</u> for 6 years or more and in <u>exposed situations</u> for up to 3 years or more.

The <u>actual period of efficacy</u> will depend on many factors such as termite hazard, climatic conditions, soil types and soil disturbance and gardening/landscaping practices.

27/04/05	
Country Micro-Lo Pre-Construction/Post Construction Termiticid	e & Insecticide

#### DIRECTIONS FOR USE: GENERAL PEST CONTROL

#### RESTRAINTS

#### **DO NOT** spray polycarbonate surfaces/roof sheeting or aged vinyl wall cladding as solvent may cause etching.

SITUATION			CRITICAL COMMENTS
Domestic and public places, commercial and industrial areas.	infestations)		Apply as a coarse, low pressure spray to the point of run-off, to cracks, crevices, harbourages, eaves, downpipes and other places where the pests may occur.
	Spiders		For optimum control of webbing spiders, use a 2-part treatment. After applying as a coarse, low pressure spray to harbourages where the spiders may occur, apply a light spray over surfaces of the building.
	Silverfish	55mL/10L	
	Cockroaches (light infestations)	of water	
	Argentine ants	least 1L spray/ 10 m <sup>2</sup> infested area	Locate ant nests and treat appropriately. Spray ant tracks or where ant activity is noticed. Apply to paths in continuous 300mm bands. Apply to base of buildings, walls, fences, rock-works, trunks of shrubs and trees, and other hard surfaces to a height of 300mm.
Domestic and public places commercial and industrial	Fleas (outdoor use only)	110mL/10L of water	Apply as a fine droplet spray. Outdoors only. Treat areas where animals frequent.
areas.	• /		Remove animals during treatment and until spray deposit is dry.
			Do not treat pets with this product. Pets should be treated with a product registered for application to animals.
Hides/Skins			Apply spray to flesh side of skins or hides sufficient to moisten them. Ensure coverage of ears and lugs. To minimise the chance of later infestations, storage area should be sprayed regularly. Repeat application every 3 months. Access through bales should be maintained for application of product.
Light vegetation	Mosquito larvae	30mL/ha	Dilute with water and apply as a spray to areas infested with mosquitoes.
Medium vegetation	-	65mL/ha	
Heavy vegetation		120mL/ha	
Light to medium vegetation	Mosquito adults	65mL/ha	
Medium to heavy vegetation		120mL/ha	
Polluted water impoundments		$2mL/10,000L$ of water or $20mL/100 \text{ m}^3$ of water.	

#### NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

### GENERAL INSTRUCTIONS

Page 13 of 16

27/04/05

#### MIXING

Half fill the spray tank with water (or creosote where applicable) and add the required amount of Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide, then add the remaining water (or creosote) with an agitator running. If using a knapsack sprayer gently shake before using. Only mix sufficient chemical for each specific application.

#### **CLEANING SPRAY EQUIPMENT**

After using Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide, empty the spray equipment completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose, and drain. To wash the system, quarter fill the tank with clean water and circulate through the pump, lines, hoses and nozzles (for knapsack sprayers spray to waste through the nozzle). Drain and repeat the washing procedure twice.

Disposal of rinsate/rinse water in accordance with Storage and Disposal instructions below.



#### INSECTICIDE RESISTANCE WARNING

For insecticide resistance management Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide is a Group 1B insecticide. Some naturally occurring insect biotypes resistant to Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide and other Group 1B insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide or other Group 1B insecticides are used repeatedly. The effectiveness of Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, A & C Rural Pty Ltd accepts no liability for any losses that may result from the failure of Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide to control resistant individuals.

Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier or A & C Rural Pty Ltd Representative.

#### PRECAUTIONS

DO NOT apply inside buildings except as a crack and crevice treatment. DO NOT apply to surface areas such as interior floors or walls.

#### **RE-ENTRY TO TREATED AREAS**

#### **Pre-construction termite management:**

Suspended Floors: DO NOT allow entry until treated areas are completely dry (normally 3-4 hours).

Concrete Slabs: Cover treated areas immediately with a moisture membrane. Re-entry permitted after laying of the moisture membrane.

#### Post-construction termite management and general pest control:

DO NOT permit re-occupation of any premises until treated areas are completely dry (normally 3-4 hours) and adequately ventilated.

#### PROTECTION OF PETS AND LIVESTOCK

Before spraying remove animals and pets from buildings and other areas to be treated. Cover or remove any open food and water containers.

Cover or remove fish tanks before spraying. DO NOT allow animals and pets to contact treated areas for at least 24 hours. Dangerous to bees. DO NOT spray any plants in flower while bees are actively foraging.

#### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

VERY HIGHLY TOXIC TO FISH AND AQUATIC INVERTEBRATES. Rinse waters and run-off from treated areas MUST NOT enter drains or waterways. For under-slab treatments the moisture membrane MUST be installed immediately after treatment. DO NOT apply to waterlogged soils. DO NOT apply if heavy rains are expected to occur within 48 hours of application.

HIGHLY TOXIC TO BIRDS. Do NOT treat fill unless it has been placed back in the trench to form the chemical soil barrier.

DO NOT spray directly on to the foliage of plants as damage to some species is possible.

#### SMALL SPILL MANAGEMENT

Wear appropriate clothing and protective equipment whilst cleaning up small spills (see **SAFETY DIRECTIONS**). Apply absorbent material such as earth, sand, cat litter or clay granules to the spill. Sweep up material and contain in a refuse vessel for disposal. If spilled inside a building, wash contaminated surfaces to deactivate the chlorpyrifos with a dilute solution of bleach (sodium hypochlorite), prepared according to the bleach label instructions. Dispose of the contaminated material in accordance with **STORAGE AND DISPOSAL** instructions below.

#### STORAGE AND DISPOSAL

Store in closed, original container in a cool well-ventilated area. Do not store for long periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling break, crush or puncture and bury empty containers in a local authority landfill. If no local landfill is available, bury the containers at a depth of 500mm or more in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. Rinsate/rinse water should be disposed of in accordance with appropriate State legislation and should preferably be sprayed on to an application site or added as up to 10% of the diluent the next time this product or another similar termiticide is used. Do not put down sewers, gutters or storm water drains. In some States wastes can only be buried at a licensed landfill.

### SAFETY DIRECTIONS

#### For General Use

Product is poisonous if absorbed by skin contact, inhaled or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Obtain an emergency supply of atropine tablets 0.6 mg. Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale vapour or spray mist. When opening the container, preparing the spray or using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield or goggles.

If product is spilt on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

For Use As A Termiticide

Product is poisonous if absorbed by skin contact, inhaled or swallowed. Repeated exposure may cause allergic disorders. Repeated minor exposure may have a cumulative poisoning effect. Obtain an emergency supply of atropine tablets 0.6 mg. Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale vapour or spray mist.

When opening the container, preparing the spray and using the prepared spray, wear chemical resistant clothing buttoned to the neck and wrist and washable hat, elbow-length PVC gloves, goggles, chemical resistant footwear and half facepiece respirator with combined dust and gas cartridge.

If clothing becomes contaminated with product or wet with spray, remove clothing immediately. If product on the skin, immediately wash area with soap and water. If product is in the eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, goggles, respirator (and if rubber wash with detergent and warm water), and contaminated clothing.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. (Ph.: 13 11 26). If swallowed, give one atropine tablet every 5 minutes until dryness of the mouth occurs - if poisoned by skin absorption or through lungs, remove any contaminated clothing, wash skin thoroughly and give atropine tablets as above. Get to a doctor or hospital quickly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

#### MATERIAL SAFETY DATA SHEET

Additional information is listed on the Material Safety Data Sheet for Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide which is available from A & C Rural Pty Ltd on request.

**CONDITIONS OF SALE:** The use of Country Micro-Lo Pre-Construction/Post-Construction Termiticide and Insecticide being beyond the control of the manufacturer no warranty expressed or implied is given by A & C Rural Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and A & C Rural Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.